

The University of New South Wales  
School of Mathematics and Statistics

**Mathematics Drop-in Centre**

**ALGEBRAIC IDENTITIES**

There are a number of algebraic identities which you need to know in order to help you solve equations and simplify expressions (by an *identity* we mean an equation involving one or more variables, which is true for all values of those variables).

- Addition of fractions:

$$\frac{w}{x} + \frac{y}{z} = \frac{wz + xy}{xz} .$$

- Square of a sum:

$$(x + y)^2 = x^2 + 2xy + y^2 .$$

- The same for a sum of three terms:

$$(x + y + z)^2 = x^2 + y^2 + z^2 + 2xy + 2xz + 2yz .$$

- Difference of two squares:

$$x^2 - y^2 = (x - y)(x + y) .$$

- Difference of two powers:

$$x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + \dots + xy^{n-2} + y^{n-1}) .$$

- Sum of two powers: if  $n$  is odd then

$$x^n + y^n = (x + y)(x^{n-1} - x^{n-2}y + \dots - xy^{n-2} + y^{n-1}) .$$

Note that there is *no* formula like this if  $n$  is even: for example,  $x^2 + y^2$  cannot be factorised in any simple way.

- The Binomial Theorem to expand  $(x + y)^n$ : this is dealt with in a separate sheet.
- Power laws and logarithm laws such as  $a^x a^y = a^{x+y}$ : these are dealt with in separate sheets.

**Comments.**

- You must be able to use these identities “in both directions”. For example, if you see  $x^2 - y^2$  you should know that it can be factorised, and if you see  $(x - y)(x + y)$  you should know that it can be expanded and simplified.
- You must be able to replace the variables in all these identities by different variables, constants or expressions. From the “difference of two squares” formula we can find, for example,

$$m^2 - n^2 = (m - n)(m + n)$$

$$x^2 - 5 = (x - \sqrt{5})(x + \sqrt{5})$$

$$9a^2 - 100b^2 = (3a - 10b)(3a + 10b)$$

$$x^6 - y^6 = (x^3 - y^3)(x^3 + y^3) .$$

- Sometimes we can use a number of these identities successively in order to give a detailed factorisation of a certain expression. For instance,

$$x^6 - y^6 = (x^3 - y^3)(x^3 + y^3) \quad (\text{difference of two squares})$$

$$= (x - y)(x^2 + xy + y^2)(x^3 + y^3)$$

(difference of 3rd powers)

$$= (x - y)(x^2 + xy + y^2)(x + y)(x^2 - xy + y^2)$$

(sum of 3rd powers).

## EXERCISES.

Please try to complete the following exercises. Remember that you **cannot** expect to understand mathematics without doing lots of practice! Please do not look at the answers before trying the questions. If you get a question wrong you should go through your working carefully, find the mistake and fix it. If there is a mistake which you cannot find, or a question which you cannot even start, please consult your tutor or the Mathematics Drop-in Centre.

### 1. Expand

- (a)  $(4x + 5y)(4x - 5y)$ ;      (b)  $(s + t)^2$ ;  
(c)  $(x - 3y + 5z)^2$ ;      (d)  $(z^3 - 4)(z^3 + 4)$ ;  
(e)  $(ab - 2cd)^2$ ;      (f)  $((a + b) - c)((a + b) + c)$ ;  
(g)  $(x + y)(x^4 - x^3y + x^2y^2 - xy^3 + y^4)$ ;  
(h)  $(a + b + c + d)^2$       (i)  $(x + y)^2 - (x - y)^2$ .

### 2. Factorise

- (a)  $x^2 + 10xy + 25y^2$ ;      (b)  $x^6 - y^4$ ;  
(c)  $x^8 - y^8$ ;      (d)  $4a^2 - 5b^2$ ;  
(e)  $x^6 - 2x^3 + 1$ ;      (f)  $z^7 - 128$  (hint:  $128 = 2^7$ ).

3. (a) Factorise  $x^4 - x^2 + 1$  by writing it as  $(x^4 + 2x^2 + 1) - 3x^2$  and using the above identities.

(b) Hence factorise  $x^{12} - 1$  into linear and quadratic factors.

### 4. The sum and difference of fractions

$$\frac{1}{x-1} - \frac{2}{x^2-1} + \frac{1}{x^3-1}$$

has a common denominator  $(x-1)(x^2-1)(x^3-1)$ , but this is not the *smallest* common denominator. By factorising the three denominators, find the smallest common denominator and hence simplify the expression.

## ANSWERS.

- (a)  $16x^2 - 25y^2$ ;  
(b)  $s^2 + 2st + t^2$ ;  
(c)  $x^2 + 9y^2 + 25z^2 - 6xy + 10xz - 30yz$ ;  
(d)  $z^6 - 16$ ;  
(e)  $a^2b^2 - 4abcd + 4c^2d^2$ ;  
(f)  $a^2 + 2ab + b^2 - c^2$ ;  
(g)  $x^5 + y^5$ ;  
(h)  $a^2 + b^2 + c^2 + d^2 + 2ab + 2ac + 2ad + 2bc + 2bd + 2cd$ ;  
(i)  $4xy$ .
- (a)  $(x + 5y)^2$ ;  
(b)  $(x^3 - y^2)(x^3 + y^2)$ ;  
(c)  $(x^4 - y^4)(x^4 + y^4)$  for a start, but hopefully you can continue and get  $(x - y)(x + y)(x^2 + y^2)(x^4 + y^4)$ ;  
(d)  $(2a - \sqrt{5}b)(2a + \sqrt{5}b)$ ;  
(e)  $(x^3 - 1)^2$ ; even better,  $(x - 1)^2(x^2 + x + 1)^2$ ;  
(f)  $(z - 2)(z^6 + 2z^5 + 4z^4 + 8z^3 + 16z^2 + 32z + 64)$ .
- (a)  $(x^2 - \sqrt{3}x + 1)(x^2 + \sqrt{3}x + 1)$ ;  
(b)  $(x - 1)(x^2 + x + 1)(x + 1)(x^2 - x + 1)(x^2 + 1)$   
 $(x^2 - \sqrt{3}x + 1)(x^2 + \sqrt{3}x + 1)$ .
- The smallest common denominator is  $(x-1)(x+1)(x^2+x+1)$  and the expression is

$$\frac{x^3 + x}{(x-1)(x+1)(x^2+x+1)}.$$