

**MATHEMATICS ENRICHMENT CLUB.<sup>1</sup>**

**Solution Sheet 9, July 23, 2013**

1. Since the sequence is arithmetic  $a_n = a_1 + (n - 1)d$  where  $d$  is the common difference. With  $a_1 = 10$  this means  $a_n = 10 + (n - 1)d$ . Now using the data  $a_{a_2} = 100$  we get

$$\begin{aligned} 100 &= 10 + (a_2 - 1)d \\ 90 &= (a_2 - 1)d \\ 90 &= (10 + (2 - 1)d)d \\ 0 &= d^2 + 10d - 90, \quad \text{so} \\ d &= \frac{-9 \pm \sqrt{81 + 4 \times 90}}{2} \\ d &= -15, 6 \quad \text{and since } d > 0, \quad d = 6. \end{aligned}$$

Now we can compute  $a_{a_{a_3}}$ , so

$$\begin{aligned} a_{a_{a_3}} &= 10 + (a_{a_3} - 1)d \\ &= 10 + ((10 + (a_3 - 1)d) - 1)d \\ &= 10 + ((10 + ((10 + (3 - 1)d) - 1)d) - 1)d \\ &= 10 + ((10 + ((10 + 2 \times 6) - 1) \times 6) - 1) \times 6 \\ &= 10 + ((10 + (22 - 1) \times 6 - 1) \times 6 \\ &= 10 + (10 + 126 - 1) \times 6 \\ &= 10 + 135 \times 6 \\ &= 820. \end{aligned}$$

2. Let's talk about two moves, one makes a number bigger, and one reduces the number by one. First, take the latter, every  $n = (n - 1) + 1$  and  $n - 1 = (n - 1) \times 1$ . To make a number bigger take instead  $n = (n - 2) + 2$  so  $2(n - 2) = 2n - 4$  which is larger than  $n$  for  $n > 4$ . Using these two moves we can make the moves

$$22 \rightarrow 40 \rightarrow 76 \rightarrow 148 \rightarrow 292 \rightarrow 580 \rightarrow 1156 \rightarrow 2308 \rightarrow 2307 \rightarrow 2306 \rightarrow \dots \rightarrow 2001$$

<sup>1</sup>Some of the problems here come from T. Gagen, Uni. of Syd. and from E. Szekeres, Macquarie Uni.

3. Note

$$\begin{aligned}
 125^{100} &= 5^{300} \\
 &= \frac{10^{300}}{2^{300}} \\
 &= \frac{10^{300}}{(1024)^{30}} \\
 &= \frac{10^{300}}{(1.024 \times 10^3)^{30}} \\
 &= \frac{10^{300}}{1.024^{30} \times 10^{90}} \\
 &= \frac{10^{210}}{1.024^{30}}.
 \end{aligned}$$

It now remains to see that  $1.024^{30} < 10$  and hence there are 210 digits. Can you show this using the binomial theorem?

4. (a) See figure 1




Figure 1: The possible paths on a  $2 \times 2$  grid. .

- (b) At each vertex ask how many paths are there from the top-left to the given vertex. In fact, it is the sum of the number of paths from the top-left to the vertex immediately to the left of the given vertex and the number of paths from the top-left to the vertex immediately above the given vertex. This means the number of paths from the top-left to each vertex follows a Pascal's Triangle pattern. So there are  $\binom{40}{20}$  paths.
- (c) This generalises easily to  $\binom{2n}{n}$ .
5. Let's label the vertices of our hexagon 1 through to 6. Then we can refer to edges as  $(xy)$  where  $x$  and  $y$  are a vertex. Now since we can just re-label the hexagon however we want, let's just consider vertex 1. There are 5 edges coming off vertex 1, and since we only have 2 colours, 3 of these edges are the same colour, let's say red. And again, since we can re-label the hexagon however we want, let's suppose these edges are (12),(13) and (14). Suppose we don't have any red triangles, so (23) must be blue to prevent  $\Delta 123$  from being red, also (34) must be blue to prevent  $\Delta 134$  from being




Figure 2: Pascals triangle emerging

all red and (24) must be blue to prevent  $\Delta 124$  from being all red. But now (23), (34) and (24) are all blue, so  $\Delta 234$  is blue.

Seeing as it doesn't matter how you label the hexagon, this always happens.

6. Look at the sequences in base-8,  $x_n$  always ends in either 3 or 5, while  $y_n$  always ends in 1. Hence they can never be the same.